ELSEVIER

Statistics & Probability Letters 64 (2003) 443–444

STATISTICS & PROBABILITY LETTERS

www.elsevier.com/locate/stapro

# Author Index for Volume 64 (2003)


| | | |
|---|---|---|
| Abry, P., *see* Pipiras, V. | (2) | 147–157 |
| Aliev, F.A., A comment on 'Unimodality of the distribution of record statistics' | (1) | 39– 40 |
| Anderson-Cook, C.M., *see* Zahran, A. | (1) | 9– 16 |
| Balakrishnan, N. and J. Mi, Existence and uniqueness of the MLEs for normal distribution based on general progressively Type-II censored samples | (4) | 407–414 |
| Balakrishnan, N., *see* Stepanov, A.V. | (1) | 69– 81 |
| Baran, R.H., *see* Jernigan, R.W. | (1) | 17– 23 |
| Bekker, L. and J. Mi, Shape and crossing properties of mean residual life functions | (3) | 225–234 |
| Belopolskaya, Y. and G.N. Milstein, An approximation method for Navier–Stokes equations based on probabilistic approach | (2) | 201–211 |
| Berrendero, J.R., Uniform strong consistency of robust estimators | (2) | 159–168 |
| Bhattacharjee, M.C., S. Ravi, R. Vasudeva and N.R. Mohan, New order preserving properties of geometric compounds | (2) | 113–120 |
| Cook, S., Modified unit root tests and momentum threshold autoregressive processes | (1) | 83– 88 |
| Csörgő, S., Rates in the complete convergence of bootstrap means | (4) | 359–368 |
| de Haan, L., *see* Drees, H. | (1) | 51– 62 |
| Delmas, C., Projections on spherical cones, maximum of Gaussian fields and Rice's method | (3) | 263–270 |
| Deng, D., On the self-normalized bounded laws of iterated logarithm in Banach space | (3) | 277–286 |
| Dindar, Z., Some more results on increments of the partially observed empirical process | (1) | 25– 37 |
| Drees, H., L. de Haan and D. Li, On large deviation for extremes | (1) | 51– 62 |
| Evangelaras, H. and C. Koukouvinos, Effects confounded with blocks in factorial designs: a projective geometric approach with two blocks | (1) | 105–111 |
| Febrero-Bande, M., *see* García-Soidán, P.H. | (2) | 169–179 |
| Fierro, R., A test of goodness of fit testing for stochastic intensities associated to counting processes | (3) | 287–292 |
| Gao, X., *see* Li, G. | (1) | 63– 68 |
| García-Soidán, P.H., W. González-Manteiga and M. Febrero-Bande, Local linear regression estimation of the variogram | (2) | 169–179 |
| Geng, Z. and K. Li, Factorization of posteriors and partial imputation algorithm for graphical models with missing data | (4) | 369–379 |
| Gilmour, S.G. and R. Mead, A Bayesian design criterion for locating the optimum point on a response surface | (3) | 235–242 |
| Gneri, M.A., Resistant estimators for stationary ergodic stochastic processes | (1) | 97–103 |
| González-Manteiga, W., *see* García-Soidán, P.H. | (2) | 169–179 |
| Grigelionis, B. and V. Mackeviius, The finiteness of moments of a stochastic exponential | (3) | 243–248 |
| Hall, P.G. and R.J. Hyndman, Improved methods for bandwidth selection when estimating ROC curves | (2) | 181–189 |
| Hettmansperger, T.P., *see* Nadar, M. | (4) | 431–442 |
| Hofmann, G., *see* Stepanov, A.V. | (1) | 69– 81 |
| Huang, M., *see* Li, G. | (1) | 63– 68 |
| Hutson, A.D., Nonparametric estimation of normal ranges given one-way ANOVA random effects assumptions | (4) | 415–424 |
| Hyndman, R.J., *see* Hall, P.G. | (2) | 181–189 |
| Janssen , A., A nonparametric Cramér–Rao inequality for estimators of statistical functionals | (4) | 347–358 |
| Jarušková , D., Asymptotic distribution of a statistic testing a change in simple linear regression with equidistant design | (1) | 89– 95 |
| Jernigan, R.W. and R.H. Baran, Testing lumpability in Markov chains | (1) | 17– 23 |
| Jurek, Z.J., Generalized Lévy stochastic areas and self decomposability | (2) | 213–222 |
| Kim, H.-M. and B.K. Mallick, A note on Bayesian spatial prediction using the elliptical distribution | (3) | 271–276 |
| Kim, J.H. and S. Lee, Tail bound for the minimal spanning tree of a complete graph | (4) | 425–430 |
| Kotz, S., *see* Yang, Z. | (4) | 335–345 |
| Koukouvinos, C., *see* Evangelaras, H. | (1) | 105–111 |
| Lee, S., *see* Kim, J.H. | (4) | 425–430 |
| Lepeltier, J.P., Nonanticipative risk sensitive control: the martingale method | (2) | 191–199 |
| Li, D., *see* Drees, H. | (1) | 51– 62 |
| Li, G., X. Gao and M. Huang, Testing multivariate one-sided hypotheses | (1) | 63– 68 |


Elsevier Science B.V.
PII: S0167-7152(03)00245-1

| | | |
|---|---|---|
| Li, H., Association of multivariate phase-type distributions, with applications to shock models | (4) | 381–392 |
| Li, K., *see* Geng, Z. | (4) | 369–379 |
| Liu, W. and W. Yang, A class of strong limit theorems for the sequences of arbitrary random variables | (2) | 121–131 |
| Mackeviius, V., *see* Grigelionis, B. | (3) | 243–248 |
| Mallick, B.K., *see* Kim, H.-M. | (3) | 271–276 |
| Maltz, A.L. and J.D. Samur, On a CLT for Gibbs fields and its functional version | (3) | 323–333 |
| Mead, R., *see* Gilmour, S.G. | (3) | 235–242 |
| Mi, J., *see* Bekker, L. | (3) | 225–234 |
| Mi, J., *see* Balakrishnan, N. | (4) | 407–414 |
| Milstein, G.N., *see* Belopolskaya, Y. | (2) | 201–211 |
| Mohan, N.R., *see* Bhattacharjee, M.C. | (2) | 113–120 |
| Muliere, P., *see* Walker, S. | (1) | 1– 7 |
| Nadar, M., T.P. Hettmansperger and H. Oja, The asymptotic covariance matrix of the Oja median | (4) | 431–442 |
| Oja, H., *see* Nadar, M. | (4) | 431–442 |
| Ottoy, J.P., *see* Thas, O. | (3) | 255–261 |
| Pipiras, V., M.S. Taqqu and P. Abry, Can continuous-time stationary stable processes have discrete linear representations? | (2) | 147–157 |
| Raqab, M.Z., P-Norm bounds for moments of progressive type II censored order statistics | (4) | 393–402 |
| Ravi, S., *see* Bhattacharjee, M.C. | (2) | 113–120 |
| Rodríguez-Lallena, J.A. and M. Úbeda-Flores, Distribution functions of multivariate copulas | (1) | 41– 50 |
| Samur, J.D., *see* Maltz, A.L. | (3) | 323–333 |
| Shapiro, A., Scheffe's method for constructing simultaneous confidence intervals subject to cone constraints | (4) | 403–406 |
| Shimizu, K. and M. Tanaka, Expected number of level-crossings for a strictly stationary ellipsoidal process | (3) | 305–310 |
| So, B.S., Maximized log-likelihood updating and model selection | (3) | 293–303 |
| Stepanov, A.V., N. Balakrishnan and G. Hofmann, Exact distribution and Fisher information of weak record values | (1) | 69– 81 |
| Tanaka, M., *see* Shimizu, K. | (3) | 305–310 |
| Taqqu, M.S., *see* Pipiras, V. | (2) | 147–157 |
| Tenreiro, C., On the asymptotic normality of multistage integrated density derivatives kernel estimators | (3) | 311–322 |
| Thas, O. and J.P. Ottoy, Some generalizations of the Anderson–Darling statistic | (3) | 255–261 |
| Úbeda-Flores, M., *see* Rodríguez-Lallena, J.A. | (1) | 41–50 |
| Vasudeva, R., *see* Bhattacharjee, M.C. | (2) | 113–120 |
| Walker, S. and P. Muliere, A bivariate Dirichlet process | (1) | 1– 7 |
| Wen, J.-W. and L.-X. Zhang, Precise asymptotics in laws of the iterated logarithm for Wiener local time | (2) | 133–145 |
| Wendl, M.C., Collision probability between sets of random variables | (3) | 249–254 |
| Yang, W., *see* Liu, W. | (2) | 121–131 |
| Yang, Z. and S. Kotz, Center-similar distributions with applications in multivariate analysis | (4) | 335–345 |
| Zahran, A. and C.M. Anderson-Cook, A general equation and optimal design for a 2-factor restricted region | (1) | 9– 16 |
| Zhang, L.-X., *see* Wen, J.-W. | (2) | 133–145 |